Docket No.: 564462001802

### Amendment to the Claims:

Please amend the claims as follows:

Please cancel claims 89 to 92, without prejudice or disclaimer

This listing of claims will replace all prior versions and listings of the claims in the application:

## **Listing of Claims:**

Claim 1 (currently amended): A recombinant expression system comprising <u>an isolated</u> [[a]] host cell comprising a nucleic acid encoding a phytase enzyme

- (i) having comprising the amino acid sequence of as set forth in SEQ ID NO:2,
- (ii) having comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432;
- (iii) having comprising the amino acid sequence of as set forth in SEQ ID NO:2 but lacking comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,
- (iv) having comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432 but lacking comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence, or
- (v) having comprising the amino acid sequence of (i), (ii), (iii) or (iv) and further comprising a sequence a coding and/or non-coding sequence imparting a desired characteristic,

wherein the nucleic acid is operably linked to a transcription control sequence.

Claim 2 (currently amended): A vector comprising a nucleic acid

- (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1,
- (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296,
- (iii) comprising the nucleotide sequence of (i) or (ii) <u>lacking further comprising</u> a <u>nucleic</u> acid encoding a homologous signal sequence or comprising a <u>nucleic acid encoding a heterologous</u> signal sequence in place of the <u>nucleic acid encoding the</u> homologous signal sequence, or

(iv) comprising the nucleotide sequence of (i), (ii), or (iii), further comprising a <u>nucleotide</u> sequence encoding a <u>coding and/or non-coding sequence polypeptide imparting a desired</u> characteristic.

Claim 3 (previously presented): The expression system of claim 1 wherein the transcription control sequence comprises a constitutive promoter.

Claim 4 (previously presented): The expression system of claim 1 wherein the transcription control sequence comprises a tissue-specific promoter.

Claim 5 (currently amended): The expression system of claim 1 wherein said <u>isolated</u> host cell is a prokaryotic cell.

Claim 6 (currently amended): The expression system of claim 1 wherein said <u>isolated</u> host cell is a eukaryotic cell.

Claim 7 (currently amended): The expression system of claim 1 wherein said <u>isolated</u> host cell is a plant cell.

Claim 8 (previously presented): The expression system of claim 1 or 16 wherein the nucleic acid further comprises a heterologous sequence encoding a signal peptide or a transit peptide.

Claim 9 (previously presented): The expression system of claim 8 wherein said signal peptide is a pathogenesis-related (PR) protein PR-S signal peptide from tobacco.

Claim 10 (currently amended): A prokaryotic cell comprising an exogenous nucleic acid encoding a phytase enzyme, wherein the nucleic acid is operably linked to a transcriptional control sequence and the phytase enzyme

(i) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1,

- (ii) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296,
- (iii) is encoded by the nucleic acid of (i) or (ii), eomprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence,
  - (iv) comprises the [[an]] amino acid sequence of as set-forth in SEQ ID NO:2,
- (v) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432,
- (vi) comprises an amino acid sequence of (iv) or (v), <u>lacking further-comprising</u> a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,
- (vii) is encoded by the nucleic acid of (i), (ii), or (iii) and further comprising a <u>nucleotide</u> sequence encoding a <u>polypeptide imparting a desired characteristic coding and/or non-coding</u> <u>sequence</u>, or
- (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a polypeptide imparting a desired characteristic coding and/or non-coding sequence.

Claim 11 (currently amended): A eukaryotic cell comprising an exogenous nucleic acid encoding a phytase enzyme, wherein the nucleic acid is operably linked to a transcriptional control sequence and the phytase enzyme

- (i) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1,
- (ii) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296,

(iii) is encoded by the nucleic acid of (i) or (ii), eomprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence,

- (iv) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2,
- (v) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432,
- (vi) comprises an amino acid sequence of (iv) or (v), <u>lacking</u> further comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,
- (vii) is encoded by the nucleic acid of (i), (ii), or (iii) and further comprising a <u>nucleotide</u> sequence encoding a <del>polypeptide imparting a desired characteristic</del> coding and/or non-coding <u>sequence</u>, or
- (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a polypeptide imparting a desired characteristic coding and/or non-coding sequence.

Claim 12 (currently amended): A cell comprising an exogenous nucleic acid encoding a phytase enzyme, wherein the nucleic acid is operably linked to a transcriptional control sequence and the phytase enzyme

- (i) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1,
- (ii) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296,
- (iii) is encoded by the nucleic acid of (i) or (ii) eomprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence,
  - (iv) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2,
- (v) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432,

(vi) comprises an amino acid of (iv) or (v) <u>lacking further comprising</u> a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,

- (vii) is encoded by the nucleic acid of (i), (ii), or (iii) and further comprising a <u>nucleotide</u> sequence encoding a <del>polypeptide imparting a desired characteristic coding and/or non-coding</del> sequence, or
- (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a polypeptide imparting a desired characteristic coding and/or non-coding sequence.

Claim 13 (previously presented): A method for making a phytase in a cell comprising culturing the cell of claim 12 under conditions wherein the phytase enzyme is expressed.

Claims 14 to 15 (canceled)

Claim 16 (currently amended): An expression system for making a polypeptide having phytase activity, comprising an isolated [[a]] host cell and an exogenous nucleic acid, wherein the exogenous nucleic acid codes for the polypeptide having phytase activity and wherein the isolated host cell is capable of expressing the polypeptide, and the polypeptide having phytase activity

- (i) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1,
- (ii) is encoded by a nucleic acid comprising the [[a]] sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296,
- (iii) is encoded by the nucleic acid of (i) or (ii) and further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence, or
- (iv) is encoded by the nucleic acid of (i), (ii) or (iii) and further comprising a <u>nucleotide</u> sequence encoding a <u>polypeptide imparting a desired characteristic coding and/or non-coding sequence</u>.

17. (previously presented): The expression system of claim 16, wherein the exogenous nucleic acid is operably linked to a transcriptional control sequence.

- 18. (previously presented): The expression system of claim 7 wherein said plant cell is a higher plant cell.
- 19. (previously presented): The expression system of claim 8 wherein the signal peptide is a secretory signal peptide.
- 20. (previously presented): The expression system of claim 1 or claim 16, wherein the nucleic acid further comprises a promoter sequence, a secretory sequence, a stabilizing sequence, a targeting sequence or a termination sequence.
- 21. (previously presented): The expression system of claim 1 or claim 16, wherein the nucleic acid is contained in a vector.
- 22. (previously presented): The expression system of claim 21, wherein the vector comprises at least a portion of a nucleotide sequence taken from a cloning vector, an expression vector, a bacterial vector, a plasmid, a viral particle, a phage, chromosomal DNA, nonchromosomal DNA, synthetic DNA, a vaccinia vector, an adenovirus vector, a fowl pox virus, a pseudorabies vector or a combination of nucleotide sequences thereof.
- 23. (previously presented): The vector of claim 2, wherein the vector comprises at least a portion of a nucleotide sequence taken from a cloning vector, an expression vector, a bacterial vector, a plasmid, a viral particle, a phage, chromosomal DNA, nonchromosomal DNA, synthetic DNA, a vaccinia vector, an adenovirus vector, a fowl pox virus, a pseudorabies vector or a combination of nucleotide sequences thereof.

24. (previously presented): The eukaryotic cell of claim 11, wherein the eukaryotic cell is a plant cell.

- 25. (previously presented): The eukaryotic cell of claim 24, wherein the plant cell is a higher plant cell.
- 26. (previously presented): The eukaryotic cell of claim 24, wherein the plant cell is a seed cell.
- 27. (previously presented): The eukaryotic cell of claim 24, wherein the plant cell is an edible flower cell, a cauliflower cell, an artichoke cell, a fruit cell, an apple cell, a banana cell, a berry cell, a currant cell, a cherry cell, a cucumber cell, a grape cell, a lemon cell, a melon cell, a nut cell, an orange cell, a peach cell, a pear cell, a plum cell, a strawberry cell, a tomato cell, a leaf cell, an alfalfa cell, a cabbage cell, an endive cell, a leek cell, a lettuce cell, a spinach cell, a tobacco cell, a root cell, an arrowroot cell, a beet cell, a carrot cell, a cassava cell, a turnip cell, a radish cell, a yam cell, a sweet potato cell, a bean cell, a pea cell, a soybean cell, a wheat cell, a barley cell, a corn cell, a rice cell, a rapeseed cell, a millet cell, a sunflower cell, an oat cell, a tuber cell, a kohlrabi cell or a potato cell.
- 28. (previously presented): The method of claim 13, further comprising converting the cell into a composition suitable for animal feed.
- 29. (previously presented): The method of claim 13, wherein the cell is a prokaryotic cell or a eukaryotic cell.
- 30. (previously presented): The method of claim 29, wherein the eukaryotic cell is a plant cell.
- 31. (previously presented): The method of claim 30, wherein the plant cell is a higher plant cell.

Application No.: 09/777,566 10 Docket No.: 564462001802

32. (previously presented): The method of claim 30, wherein the plant cell is a seed cell.

33. (previously presented): The method of claim 30 wherein the plant cell is an edible flower cell, a cauliflower cell, an artichoke cell, a fruit cell, an apple cell, a banana cell, a berry cell, a currant cell, a cherry cell, a cucumber cell, a grape cell, a lemon cell, a melon cell, a nut cell, an orange cell, a peach cell, a pear cell, a plum cell, a strawberry cell, a tomato cell, a leaf cell, an alfalfa cell, a cabbage cell, an endive cell, a leek cell, a lettuce cell, a spinach cell, a tobacco cell, a root cell, an arrowroot cell, a beet cell, a carrot cell, a cassava cell, a turnip cell, a radish cell, a yam cell, a sweet potato cell, a bean cell, a pea cell, a soybean cell, a wheat cell, a barley cell, a corn cell, a rice cell, a rapeseed cell, a millet cell, a sunflower cell, an oat cell, a tuber cell, a kohlrabi cell or a potato cell.

#### 34. (canceled)

- 35. (currently amended): A vector comprising a nucleic acid
- (i) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (ii) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432,
- (iii) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 and lacking comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence;
- (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432 and <u>lacking comprising</u> a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,
- (v) encoding a polypeptide comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a sequence encoding a polypeptide imparting a desired characteristic coding and/or non-coding sequence, or

Docket No.: 564462001802

(vi) a nucleic acid completely complementary to any of the nucleic acids of [[steps]] (i) to(v),

wherein optionally the signal sequence comprises a secretory signal peptide.

- 36. (currently amended): A cell comprising a vector comprising a nucleic acid
- (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1;
- (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296,
- (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence,
- (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; or,
- (vi) encoding the polypeptide of (iv) or (v) <u>lacking</u> further comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,
- (vii) comprising the nucleic acid of (i), (ii), or (iii) and further comprising a nucleotide sequence encoding a coding and/or non-coding sequence,
- (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a coding and/or non-coding sequence, or
- ([[vi]]ix) comprising a sequence that is the complete complement of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

wherein optionally the signal sequence comprises a secretory signal peptide.

# 37. (canceled)

38. (currently amended): A cloning vector comprising a nucleic acid (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprising the [[a]] nucleotide sequence of as set-forth in SEQ ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence, (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (vi) encoding the polypeptide of (iv) or (v) lacking further eomprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; (vii) comprising the nucleic acid of (i), (ii), or (iii) and further comprising a nucleotide sequence encoding a coding and/or non-coding sequence; (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a coding and/or non-coding sequence; or, ([[vii]]ix) comprising a sequence that is the complete complement of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

wherein optionally the signal sequence comprises a secretory signal peptide.

the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence, (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (vi) encoding the polypeptide of (iv) or (v) lacking further comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; (vii) comprising the nucleic acid of (i), (ii), or (iii) and further

comprising a nucleotide sequence encoding a coding and/or non-coding sequence; (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a coding and/or non-coding sequence; or, ([[vii]]ix) comprising a sequence that is the complete complement of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

wherein optionally the signal sequence comprises a secretory signal peptide.

40. (currently amended): A bacterial vector comprising a nucleic acid (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) further-comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence, (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (vi) encoding the polypeptide of (iv) or (v) lacking further comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; (vii) comprising the nucleic acid of (i), (ii), or (iii) and further comprising a nucleotide sequence encoding a coding and/or non-coding sequence; (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a coding and/or non-coding sequence; or, ([[vii]]ix) comprising a sequence that is the complete complement of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

wherein optionally the signal sequence comprises a secretory signal peptide.

41. (currently amended): A plasmid comprising a nucleic acid (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal

Application No.: 09/777,566

sequence or comprising a <u>nucleic acid sequence encoding a</u> heterologous signal sequence in place of the <u>nucleic acid sequence encoding the</u> homologous signal sequence, (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (vi) encoding the polypeptide of (iv) or (v) <u>lacking further comprising</u> a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; (vii) comprising the nucleic acid of (i), (ii), or (iii) and further comprising a nucleotide sequence encoding a coding and/or non-coding sequence; (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a coding and/or non-coding sequence; or, ([[vii]]ix) comprising a sequence that is the complete complement of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

14

Docket No.: 564462001802

wherein optionally the signal sequence comprises a secretory signal peptide.

42. (currently amended): A viral particle comprising a nucleic acid (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence, (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (vi) encoding the polypeptide of (iv) or (v) lacking further comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; (vii) comprising the nucleic acid of (i), (ii), or (iii) and further comprising a nucleotide sequence encoding a coding and/or non-coding sequence; (viii) comprises the amino acid sequence; or, ([[vii]]ix) comprising a sequence that is the complete complement

of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

wherein optionally the signal sequence comprises a secretory signal peptide.

43. (currently amended): A phage comprising a nucleic acid (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprising the [[a]] nucleotide sequence of as set forth-in SEO ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid sequence encoding a heterologous signal sequence in place of the nucleic acid sequence encoding the homologous signal sequence, (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set-forth in SEQ ID NO:2 from amino acid residue 1 to 432; (vi) encoding the polypeptide of (iv) or (v) lacking further comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; (vii) comprising the nucleic acid of (i), (ii), or (iii) and further comprising a nucleotide sequence encoding a coding and/or non-coding sequence; (viii) comprises the amino acid sequence of (iv), (v) or (vi) and further comprising a sequence encoding a coding and/or non-coding sequence; or, ([[vii]]ix) comprising a sequence that is the complete complement of (i), (ii), [[or]] (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), [[or]] (vi), or (viii),

wherein optionally the signal sequence comprises a secretory signal peptide.

### 44. (canceled)

45. (currently amended): A recombinant expression system comprising a nucleic acid encoding a phytase enzyme (i) having the amino acid sequence of as set forth in SEQ ID NO:2 comprising lacking a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence, or (ii) having the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432 lacking comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence,

wherein optionally the signal sequence comprises a secretory signal peptide.

46. (currently amended): A method for making a phytase in a cell, wherein the phytase is encoded by an exogenous nucleic acid, comprising culturing the cell under conditions wherein the phytase is expressed,

wherein the exogenous nucleic acid (i) comprises the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1; (ii) comprises the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296, (iii) comprising the nucleotide sequence of (i) or (ii) lacking a nucleic acid sequence encoding further comprising a homologous signal sequence or comprising a nucleic acid encoding a heterologous signal sequence in place of the nucleic acid encoding a homologous signal sequence; or (iv) comprising a nucleotide sequence as set forth in (i), (ii) or (iii) but lacking a signal sequence, or (v) comprising the [[a]] nucleotide sequence of as set forth in (i), (ii), or (iii) or (iii) or (iv) and a heterologous nucleotide sequence encoding a imparting a desired eharacteristic coding and/or non-coding sequence,

wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 47 to 80 (canceled)

Claim 81 (currently amended): A method for making a phytase in a cell, wherein the phytase is encoded by an exogenous nucleic acid, comprising culturing the cell under conditions wherein the phytase is expressed,

wherein the exogenous nucleic acid (i) encodes a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2, (ii) encodes a polypeptide comprising the [[an]] amino acid of as set forth in SEQ ID NO:2 from amino acid residues 1 to 432, (iii) encodes the polypeptide of (i) or (ii) lacking comprising a homologous signal sequence or a heterologous signal sequence in place of the homologous signal sequence, or (iv) encoding a polypeptide comprising an amino acid sequence as set forth in (i), (ii) or (iii) but lacking a signal sequence, or (v) encoding a polypeptide comprising the [[a]] amino acid sequence of as set forth in (i), (ii), or (iii) or (iv) and a heterologous coding and/or non-coding amino acid sequence imparting a desired characteristic,

wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 82 (currently amended): An expression system for making a polypeptide having phytase activity, comprising <u>an isolated</u> [[a]] host cell and an exogenous nucleic acid, wherein the exogenous nucleic acid codes for the polypeptide having phytase activity and wherein the <u>isolated</u> host cell is capable of expressing the polypeptide,

and the polypeptide having phytase activity (i) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2, (ii) comprises the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432, (iii) comprises the amino acid sequence of (i) or (ii), wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence, (iv) comprises the amino acid sequence of (i), (ii) or (iii) and further comprising comprises a sequence encoding a polypeptide imparting a desired characteristic coding and/or non-coding sequence, or (v) comprises enzymatically active fragments of (i), (ii), (iii), or (iv),

wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 83 (currently amended): A recombinant expression system comprising an isolated [[a]] host cell comprising a nucleic acid

(1) [[(i)]] encoding a phytase (i) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (ii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (iii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2, wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; [[or]] (iv) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432, wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence, or (v) comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a coding and/or non-coding sequence

wherein the phytase sequence of (i), (ii), (iii), [[or]] (iv), or (v) has at least one conservative amino acid substitution from SEQ ID NO:2, and the conservative amino acid substitution comprises (a) a replacement, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile, (b) an interchange of the hydroxyl residues Ser and Thr, (c) an exchange of the acidic residues Asp and Glu, (d) a substitution between the amide residues Asn and Gln, (e) an exchange of the basic residues Lys and Arg, (f) a replacement among the aromatic residues Phe, Tyr, or (g) any combination of a, b, c, d, e or f; or

(2) a nucleic acid comprising a sequence that is the complete complement of (1); wherein the nucleic acid is operably linked to a transcriptional control sequence, and the amino acid sequence of the phytase has at least 95% sequence identity to SEQ ID NO:2, wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 84 (currently amended): A recombinant expression system comprising an isolated [[a]] host cell comprising

(1) a nucleic acid encoding an enzymatically active fragment of a phytase, wherein the phytase comprises the [[an]] amino acid sequence of as set forth in (i) SEQ ID NO:2; (ii) residues 1 to 432 of SEQ ID NO:2; (iii) SEQ ID NO:2 lacking comprising a homologous signal sequence or comprising a heterologous signal sequence in place of the homologous signal sequence; [[or]] (iv) SEQ ID NO:2 from amino acid residue 1 to 432, wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; or (v) comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a coding and/or non-coding sequence,

wherein the phytase sequence of (i), (ii), (iii), [[or]] (iv), or (v) has at least one conservative amino acid substitution from SEQ ID NO:2, wherein the conservative amino acid substitution comprises (a) a replacement, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile, (b) an interchange of the hydroxyl residues Ser and Thr, (c) an exchange of the acidic residues Asp and Glu, (d) a substitution between the amide residues Asn and Gln, (e) an exchange of the basic residues Lys and Arg, (f) a replacement among the aromatic residues Phe, Tyr, or (g) any combination of a, b, c, d, e or f, or

Docket No.: 564462001802

(2) a nucleic acid comprising a sequence that is the complete complement of (1); wherein the nucleic acid is operably linked to a transcriptional control sequence, and the amino acid sequence of the phytase has at least 95% sequence identity to SEQ ID NO:2, wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 85 (currently amended): A vector comprising a nucleic acid

(1) [[(i)]] encoding a phytase (i) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (ii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (iii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2, wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; [[or]] (iv) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432, wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; or (v) comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a coding and/or non-coding sequence,

wherein the phytase sequence of (i), (ii), (iii), [[or]] (iv), or (v) has at least one conservative amino acid substitution from SEQ ID NO:2, and the conservative amino acid substitution comprises (a) a replacement, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile, (b) an interchange of the hydroxyl residues Ser and Thr, (c) an exchange of the acidic residues Asp and Glu, (d) a substitution between the amide residues Asn and Gln, (e) an exchange of the basic residues Lys and Arg, (f) a replacement among the aromatic residues Phe, Tyr, or (g) any combination of a, b, c, d, e or f; or

(2) a nucleic acid comprising a sequence that is the complete complement of (1); wherein the nucleic acid is operably linked to a transcriptional control sequence, and the amino acid sequence of the phytase has at least 95% sequence identity to SEQ ID NO:2,

wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 86 (currently amended): An isolated [[A]] host cell comprising a vector comprising a nucleic acid

(1) [[(i)]] encoding a phytase (i) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2; (ii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432; (iii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2, wherein the amino acid sequence lacks comprising a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; [[or]] (iv) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432, wherein the amino acid sequence lacks comprises a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; or (v) comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a coding and/or non-coding sequence,

wherein the phytase sequence of (i), (ii), (iii), [[or]] (iv), or (v) has at least one conservative amino acid substitution from SEQ ID NO:2, and the conservative amino acid substitution comprises (a) a replacement, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile, (b) an interchange of the hydroxyl residues Ser and Thr, (c) an exchange of the acidic residues Asp and Glu, (d) a substitution between the amide residues Asn and Gln, (e) an exchange of the basic residues Lys and Arg, (f) a replacement among the aromatic residues Phe, Tyr, or (g) any combination of a, b, c, d, e or f; or

(2) a nucleic acid comprising a sequence that is the complete complement of (1); wherein the nucleic acid is operably linked to a transcriptional control sequence, and the amino acid sequence of the phytase has at least 95% sequence identity to SEQ ID NO:2,

wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 87 (currently amended): An isolated [[A]] cell comprising an exogenous nucleic acid
(1) [[(i)]] encoding a phytase (i) comprising the [[an]] amino acid sequence of as set forth in
SEQ ID NO:2; (ii) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from
amino acid residue 1 to 432; (iii) comprising the [[an]] amino acid sequence of as set forth in SEQ
ID NO:2, wherein the amino acid sequence lacks comprising a homologous signal sequence or

comprises a heterologous signal sequence in place of the homologous signal sequence; [[or]] (iv) comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432, wherein the amino acid sequence lacks comprises a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; or (v) comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a coding and/or non-coding sequence,

wherein the phytase sequence of (i), (ii), (iii), [[or]] (iv), or (v) has at least one conservative amino acid substitution from SEQ ID NO:2, and the conservative amino acid substitution comprises (a) a replacement, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile, (b) an interchange of the hydroxyl residues Ser and Thr, (c) an exchange of the acidic residues Asp and Glu, (d) a substitution between the amide residues Asn and Gln, (e) an exchange of the basic residues Lys and Arg, (f) a replacement among the aromatic residues Phe, Tyr, or (g) any combination of a, b, c, d, e or f; or

(2) a nucleic acid comprising a sequence that is the complete complement of (1); wherein the nucleic acid is operably linked to a transcriptional control sequence, and the amino acid sequence of the phytase has at least 95% sequence identity to SEQ ID NO:2,

wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 88 (currently amended): An isolated [[A]] cell comprising

(1) an exogenous nucleic acid encoding an enzymatically active fragment of a phytase, wherein the phytase comprises the [[an]] amino acid sequence of as set forth in (i) SEQ ID NO:2; (ii) residues 1 to 432 of SEQ ID NO:2; (iii) SEQ ID NO:2, wherein the amino acid sequence lacks emprises a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; [[or]] (iv) SEQ ID NO:2 from amino acid residue 1 to 432, wherein the amino acid sequence lacks emprises a homologous signal sequence or comprises a heterologous signal sequence in place of the homologous signal sequence; or (v) comprising the amino acid sequence of (i), (ii), (iii), or (iv), and further comprising a coding and/or non-coding sequence,

wherein the phytase sequence of (i), (ii), (iii), [[or]] (iv), or (v) has at least one conservative amino acid substitution from SEQ ID NO:2, wherein the conservative amino acid substitution comprises (a) a replacement, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile, (b) an interchange of the hydroxyl residues Ser and Thr, (c) an exchange of the acidic residues Asp and Glu, (d) a substitution between the amide residues Asn and Gln, (e) an exchange of the basic residues Lys and Arg, (f) a replacement among the aromatic residues Phe, Tyr, or (g) any combination of a, b, c, d, e or f; or

(2) a nucleic acid comprising a sequence that is the complete complement of (1); wherein the nucleic acid is operably linked to a transcriptional control sequence, and the amino acid sequence of the phytase has at least 95% sequence identity to SEQ ID NO:2, wherein optionally the signal sequence comprises a secretory signal peptide.

Claims 89 through 91 (canceled):

Claim 92 (currently amended): An expression system comprising a nucleic acid:

- (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1;
- (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296;
- (iii) comprising the nucleotide sequence of (i) or (ii) <u>but lacking further comprising</u> a <u>nucleic</u> acid encoding a homologous signal sequence or <u>comprising</u> a <u>nucleic acid encoding a</u> heterologous signal sequence in place of the nucleic acid encoding the homologous signal sequence;
- (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432;
- (vi) encoding the polypeptide of (iv) or (v) <u>but lacking further comprising</u> a homologous signal sequence or <u>comprising</u> a heterologous signal sequence in place of the homologous signal sequence;

(vii) comprising a sequence that is the complete complement of (i), (ii) or (iii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v) or (vi);

- (viii) comprising a nucleotide sequence as set forth in (i), (ii), or (iii) but lacking a leader sequence;
- (ix) encoding a polypeptide comprising an amino acid sequence as set forth in (iv), (v) or (vi) but lacking a leader sequence;
- (x) comprising the [[a]] nucleotide sequence of as set forth in (i), (ii), or (iii) or (viii) and a heterologous nucleotide sequence encoding a imparting a desired characteristic coding and/or non-coding sequence; [[or]]
- ([[x]]viii) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in (iv), (v), or (vi) or (x) and a heterologous amino acid coding and/or non-coding sequence imparting a desired characteristic, or
- (ix) comprising a sequence that is the complete complement of (i), (ii), (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), (vi), or (viii), wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 93 (currently amended): An isolated, synthetic or recombinant nucleic acid

- (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1;
- (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296;
- (iii) comprising the nucleotide sequence of (i) or (ii) <u>but lacking further comprising</u> a <u>nucleic</u> acid encoding a homologous signal sequence or <u>comprising</u> a <u>nucleic acid encoding a</u> heterologous signal sequence in place of the <u>nucleic acid encoding the</u> homologous signal sequence;
- (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432;

- (vi) encoding the polypeptide of (iv) or (v) <u>but lacking further comprising</u> a homologous signal sequence or <u>comprising</u> a heterologous signal sequence in place of the homologous signal sequence;
- (vii) comprising a sequence that is the complete complement of (i), (ii) or (iii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v) or (vi);
- (viii) comprising a nucleotide sequence as set forth in (i), (ii), or (iii) but lacking a leader sequence;
- (ix) encoding a polypeptide comprising an amino acid sequence as set forth in (iv), (v) or (vi) but lacking a leader sequence;
- (x) comprising the [[a]] nucleotide sequence of as set forth in (i), (ii), or (iii) or (viii) and a nucleic acid encoding a a heterologous nucleotide sequence imparting a desired characteristic coding and/or non-coding sequence; [[or]]
- ([[xi]]<u>viii</u>) encoding a polypeptide comprising <u>the</u> [[an]] amino acid sequence <u>of as set forth</u> in (iv), (v), <u>or</u> (vi) <del>or</del> (vi) <del>or</del> (x) and a heterologous <del>amino acid</del> <u>coding and/or non-coding sequence</u> sequence <u>imparting a desired characteristic</u>, <u>or</u>
- (ix) comprising a sequence that is the complete complement of (i), (ii), (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), (vi), or (viii); wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 94 (currently amended): A method for glycosylating a polypeptide comprising:

- a) providing a nucleic acid comprising a sequence encoding a polypeptide having phytase activity to a cell, wherein the nucleic acid sequence
  - (i) comprises the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1;
- (ii) comprises the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296;
- (iii) comprises the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid encoding a heterologous signal sequence in place of the nucleic acid encoding the homologous signal sequence;

- (iv) encodes a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (v) encodes a polypeptide comprising the [[an]] amino acid sequence of as set-forth in SEQ ID NO:2 from amino acid residue 1 to 432;
- (vi) encodes the polypeptide of (iv) or (v) <u>but lacking further comprising</u> a homologous signal sequence or <u>comprising</u> a heterologous signal sequence in place of the homologous signal sequence;
- (vii) comprises a sequence that is the complete complement of (i), (ii) or (iii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v) or (vi);
- (viii) comprising a nucleotide sequence as set forth in (i), (ii), or (iii) but lacking a leader sequence;
- (ix) encoding a polypeptide comprising an amino acid sequence as set forth in (iv), (v) or (vi) but lacking a leader sequence;
- (x) comprising the [[a]] nucleotide sequence of as set forth in (i), (ii), or (iii) or (viii) and a heterologous nucleotide sequence encoding imparting a desired characteristic a coding and/or non-coding sequence; [[or]]
- ([[xi]]<u>viii</u>) encoding a polypeptide comprising <u>the</u> [[an]] amino acid sequence <u>of as set forth</u> in (iv), (v), <u>or</u> (vi) <del>or</del> (vi) <del>or</del> (x) and a heterologous <u>amino acid coding and/or non-coding sequence</u> sequence <u>imparting a desired characteristic; or</u>
- (ix) comprises a sequence that is the complete complement of (i), (ii), (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), (vi) or (viii), wherein optionally the signal sequence comprises a secretory signal peptide,
- b) expressing the polypeptide in the cell wherein the cell is capable of glycosylating the polypeptide.
  - Claim 95 (previously presented): The method of claim 94, wherein the cell is a yeast cell.

Claim 96 (currently amended): The method of claim 89 or claim 94, wherein the heterologous <u>nucleotide sequence of (vii)</u> or the heterologous <u>amino acid</u> sequence <u>of (viii)</u> imparting a desired characteristic comprises a secretory signal sequence.

Claim 97 (currently amended): An isolated [[A]] transformed cell or an isolated [[a]] host cell comprising an exogenous nucleic acid

- (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1;
- (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296;
- (iii) comprising the nucleotide sequence of (i) or (ii) <u>further comprising lacking a nucleic</u> acid encoding a homologous signal sequence or <u>comprising a nucleic acid encoding a heterologous</u> signal sequence in place of the <u>nucleic acid encoding the homologous signal sequence</u>;
- (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432;
- (vi) encoding the polypeptide of (iv) or (v) <u>but lacking further comprising</u> a homologous signal sequence or <u>comprising</u> a heterologous signal sequence in place of the homologous signal sequence;
- (vii) comprising a sequence that is the complete complement of (i), (ii) or (iii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v) or (vi);
- (viii) comprising a nucleotide sequence as set forth in (i), (ii), or (iii) but lacking a leader sequence;
- (ix) encoding a polypeptide comprising an amino acid sequence as set forth in (iv), (v) or (vi) but-lacking a leader sequence;
- (x) comprising the [[a]] nucleotide sequence of as set-forth in (i), (ii), or (iii) or (viii) and a heterologous nucleotide sequence encoding a imparting a desired characteristic coding and/or non-coding sequence; [[or]]

([[xi]]<u>viii</u>) encoding a polypeptide comprising an amino acid sequence <u>of</u> as set forth in (iv), (v), <u>or</u> (vi) <del>or</del> (x) and a heterologous amino acid <u>coding</u> and/or <u>non-coding</u> sequence <u>imparting</u> a desired characteristic; <u>or</u>

(ix) comprising a sequence that is the complete complement of (i), (ii), (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), (vi), or (viii), wherein optionally the signal sequence comprises a secretory signal peptide.

Claim 98 (currently amended): A vector comprising an exogenous nucleic acid

- (i) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1;
- (ii) comprising the [[a]] nucleotide sequence of as set forth in SEQ ID NO:1 from nucleotide residue 1 to 1296;
- (iii) comprising the nucleotide sequence of (i) or (ii) further comprising lacking a nucleic acid encoding a homologous signal sequence or comprising a nucleic acid encoding a heterologous signal sequence in place of the nucleic acid encoding the homologous signal sequence;
- (iv) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2;
- (v) encoding a polypeptide comprising the [[an]] amino acid sequence of as set forth in SEQ ID NO:2 from amino acid residue 1 to 432;
- (vi) encoding the polypeptide of (iv) or (v) <u>but lacking further comprising</u> a homologous signal sequence or <u>comprising</u> a heterologous signal sequence in place of the homologous signal sequence;
- (vii) comprising a sequence that is the complete complement of (i), (ii) or (iii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v) or (vi);
- (viii) comprising the [[a]] nucleotide sequence of as set forth in (i), (ii), or (iii) but lacking a leader-sequence;
- (ix) encoding a polypeptide comprising an amino acid sequence of as set forth in (iv), (v) or (vi) but lacking a leader sequence;

Docket No.: 564462001802

(x)-comprising the [[a]] nucleotide sequence of as-set forth in (i), (ii)[[,]] or (iii) [[or (viii)]] and a heterologous nucleotide sequence encoding a imparting a desired characteristic coding and/or non-coding sequence; [[or]]

([[xi]]<u>viii</u>) encoding a polypeptide comprising an amino acid sequence <u>of as set forth in</u> (iv), (vi) or (x) and a heterologous <u>amino acid coding and/or non-coding sequence imparting a desired characteristic; or</u>

(ix) comprising a sequence that is the complete complement of (i), (ii), (iii), or (vii) or the complete complement of the nucleic acid encoding the polypeptide of (iv), (v), (vi), or (viii), wherein optionally the signal sequence comprises a secretory signal peptide.